Protein Molecular Weight Marker (Broad)

Code No. 3452 **Size: for 200 lanes**
Conc.: 18 μg/μl

Supplied Reagents:

5X Loading Buffer	**1 ml**
1 M DTT (Dithiothreitol)	**100 μl**

Description :
This marker consists of 9 kinds of proteins with a Molecular Weight range between 6.5 kDa and 200 kDa. Each protein is proportioned to yield uniform band intensies when stained with Coomassie Brilliant Blue R-250 after run on SDS-polyacrylamide gel.
5 μl of 20 - fold diluted marker is loaded per lane of SDS-PAGE minigel for standard use. In this case, this marker is sufficient for 200 lanes.

Form :

50 mM	Tris-HCl, pH 6.8
1 mM	EDTA
200 mM	NaCl
50%	Glycerol

Component Proteins :

Protein	Source	MW (Da)
Myosin	Pig	200,000
β -galactosidase	*E. coli*	116,000
Phosphorylase B	Rabbit muscle	97,200
Serum Albumin	Bovine	66,409
Ovalbumin	Hen egg White	44,287
Carbonic anhydrase	Bovine	29,000
Trypsin inhibitor	Soybean	20,100
Lysozyme	Hen egg white	14,300
Aprotinin	Bovine pancreas	6,500

Storage :

Protein MW Marker (Broad), 1 M DTT : -20℃
5X Loading Buffer : Store at RT after used

5X Loading Buffer (Store at RT after used) :

250 mM	Tris-HCl, pH 6.8
6%	SDS
0.05%	BPB
50%	Glycerol

Application Example :

1. Combine the following reagents in a tube.

1 M DTT	2 μl
5X Loading Buffer	20 μl

2. Prepare 20 - fold diluted marker by adding the following components to the solution prepared in 1.

Protein MW Marker (Broad)	5 μl
Sterile purified water	73 μl

   ＊ 20 - fold diluted marker is stable for 2 - 3 months at -20℃. It is recommended to store in aliquots for several uses, not to repeat freeze-thaw cycles.

3. Mix 20 - fold diluted marker well, and heat at 100℃ for 5 minutes.
   Load 5 μl per lane of SDS-PAGE minigel.
   Run 5 - 20% gradient SDS-polyacrylamide gel electrophoresis.

4. Perform staining with Coomassie Brilliant Blue R-250.

Electrophoresis Result (5 - 20% gradient SDS-PAGE)



Note :
It is recommended to use 5 - 20% gradient SDS-polyacrylamide gel for electrophoresis.
If the concentration of gel is too low, the mobility of bands of low molecular weight is faster than the mobility of BPB. If the concentration of gel is too high, high molecular weight bands may not separate well or they may not migrate.

Note
This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals. Also, do not use this product as food, cosmetic, or household item, etc.
Takara products may not be resold or transferred, modified for resale or transfer, or used to manufacture commercial products without written approval from Takara Bio Inc.
If you require licenses for other use, please contact us from our website at www.takarabio.com.
Your use of this product is also subject to compliance with any applicable licensing requirements described on the product web page. It is your responsibility to review, understand and adhere to any restrictions imposed by such statements.
All trademarks are the property of their respective owners. Certain trademarks may not be registered in all jurisdictions.

v202510w

Protein Molecular Weight Marker (Broad)

Code No. 3452　　**容量：　200 回分**
濃度：　18 μg/μl

添付試薬：

5 × Loading Buffer	**1 ml**
1 M DTT (Dithiothreitol)	**100 μl**

●製品説明
本製品は、6.5 kDa から 200 kDa までの 9 種類の異なる分子量のタンパク質からなる分子量マーカーである。SDS-PAGE で泳動後、Coomassie Brilliant Blue R-250 で染色を行った場合、各バンドの濃さは同じになる。1 回の使用量は、SDS-PAGE ミニゲル 1 レーンあたり、本製品の 20 倍希釈液 5 μl である。その場合、本製品は約 200 回使用できる。

●形状

50 mM	Tris-HCl（pH6.8）
1 mM	EDTA
200 mM	NaCl
50%	Glycerol

●製品中に含まれるタンパク質の種類

タンパク質の種類	由来	MW (Da)
Myosin	Pig	200,000
β -galactosidase	*E. coli*	116,000
Phosphorylase B	Rabbit muscle	97,200
Serum Albumin	Bovine	66,409
Ovalbumin	Hen egg White	44,287
Carbonic anhydrase	Bovine	29,000
Trypsin inhibitor	Soybean	20,100
Lysozyme	Hen egg white	14,300
Aprotinin	Bovine pancreas	6,500

●保存

Protein MW Marker (Broad)、1 M DTT：　－ 20℃
5 × Loading Buffer：　開封後、室温保存

●5 × Loading Buffer（開封後、室温保存）

250 mM	Tris-HCl（pH6.8）
6%	SDS
0.05%	BPB
50%	Glycerol

●使用方法
1. 下記の方法で希釈用バッファーを調製する。

1 M DTT	2 μl
5 × Loading Buffer	20 μl

2. 上記で調製した溶液（22 μl）に、下記の溶液を加え、Protein MW Marker の 20 倍希釈液を調製する。

Protein MW Marker (Broad)	5 μl
滅菌精製水	73 μl

※ 調製した 20 倍希釈液は、－ 20℃で 2 ～ 3 ヵ月保存可能。必要に応じて小分け分注し、凍結融解の繰り返しは避けること。

3. 均一に混合した後、100℃で 5 分間加熱処理し、5 μl を用いて 5 ～ 20% グラジエント SDS-PAGE で電気泳動を行う。（SDS-PAGE ミニゲルの場合）

4. 泳動後、Coomassie Brilliant Blue R-250 で染色を行う。

電気泳動写真（5 ～ 20% グラジエント SDS-PAGE）



●使用上の注意
泳動には、5 ～ 20% グラジエントポリアクリルアミドゲルの使用をお勧めします。ゲル濃度が低すぎる場合、低分子量のバンドの移動速度が BPB より速くなります。ゲル濃度が高すぎる場合、高分子量のバンドの分離能が悪くなったり、泳動されない場合があります。

●注意
本製品は研究用として販売しております。ヒト、動物への医療、臨床診断用には使用しないようご注意ください。また、食品、化粧品、家庭用品等として使用しないでください。
タカラバイオの承認を得ずに製品の再販・譲渡、再販・譲渡のための改変、商用製品の製造に使用することは禁止されています。
ライセンスに関する情報は弊社ウェブカタログをご覧ください。
本データシートに記載されている会社名および商品名などは、各社の商号、または登録済みもしくは未登録の商標であり、これらは各所有者に帰属します。

v202510w

タカラバイオ株式会社
ウェブサイト　https://www.takara-bio.co.jp

製品についての技術的なお問い合わせ先
テクニカルサポートライン
Tel　077-565-6999
Fax　077-565-6995