

FITC-Labeled Monoclonal Anti-FMC63 Antibody, Mouse IgG1 (Y45)

Catalog # FM3-FY45



Source

Monoclonal Anti-FMC63 Antibody, Mouse IgG1 (Y45) is a neutralizing mouse monoclonal antibody produced from hybridoma.

Application

Flow Cytometry (Evaluation of Anti-CD19 (FMC63 scFv) CAR Expression).

Clone

Y45

Species

Mouse

Isotype

Mouse IgG1 | Mouse Kappa

Specificity

Specifically recognizes the antigen-recognition domain of FMC63 derived CARs.

Immunogen

Recombinant FMC63 scFv derived from HEK293 cells

Conjugate

FITC

Excitation source: 488 nm spectral line, argon-ion laser

Excitation Wavelength: 488 nm

Emission Wavelength: 535 nm

Isotype Control

The Isotype control is sold separately and you can search for Cat. No. [DNP-FMIAI](#) for product information.

Recommended Dilution

1:50

Formulation

Lyophilized from 0.22 μm filtered solution in PBS, 0.5% BSA, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please protect from light and avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- 20°C to -70°C for 24 months in lyophilized state;
- 70°C for 12 months under sterile conditions after reconstitution.

Bioactivity-FACS




2e5 of Anti-CD19 CAR-293 cells were stained with 100 μL of 1:50 dilution (2 μL stock solution in 100 μL FACS buffer) FITC-Labeled Monoclonal Anti-

A



B



Non-specificity of FITC-Labeled Monoclonal Anti-FMC63 Antibody, Mouse IgG1 (Y45) (Cat. No. FM3-FY45) binding to CD3+ cells present in human

Discounts, Gifts,
and more!



FITC-Labeled Monoclonal Anti-FMC63 Antibody, Mouse IgG1 (Y45)

Catalog # FM3-FY45



FMC63 Antibody, Mouse IgG1 (Y45) (Cat. No. FM3-FY45) and isotype control respectively. FITC signal was used to evaluate the binding activity (QC tested).

PBMC. Human PBMCs were simultaneously stained with PE-labeled anti-CD3 antibody and FITC-Labeled Monoclonal Anti-FMC63 Antibody, Mouse IgG1 (Y45) (2 μ L of the antibody stock solution corresponds to labeling of 5e5 cells in a final volume of 100 μ L), washed and then analyzed with FACS. Both PE and FITC positive signals was used to evaluate the non-specific binding activity to human CD3+ cells (QC tested).

Background

FMC63 is an IgG2a mouse monoclonal antibody specific for CD19, which is a target for the immunotherapy of B lineage leukaemias and lymphomas. FMC63 scFv is the most commonly used ectodomain component of CD19-specific CARs. So far, most of reported CART19 trials contain the anti-CD19 scFv derived from FMC63, including the two FDA-approved CARs Kymriah and Yescarta.

